



International Congress on Biological, Physical And Chemical Studies

International Congress on Biological, Physical And Chemical Studies - is an international conference platform under open access policy. The conference is led by international expert members who take an objective approach to peer review, ensuring each research paper is reviewed, edited by authors and evaluated on its own scholarly merits and research integration. Publishing and joining on the proceeding of the International Congress on Biological, Physical And Chemical Studies will ensure publishing experience and indexing possibilities on various global indexing.

Pregnancy and Delivery Outcomes in Women Exposed to Tobacco Smoke

Klimashkin Alexey Alexandrovich

TashPMI, Assistant Professor of the Department of Obstetrics and
Gynecology, Pediatric Gynecology, PhD

Khamdamova Nodira Ikrom kizi

Tashkent Pediatric Medical Institute

ABSTRACT

This thesis examines the effect that passive smoking has on pregnant women childbirth and neonates. Exposure to tobacco smoke, a complex of thousands of different chemicals, is dangerous for reproductive health. Carbon monoxide and nicotinic acid harm the hormonal background necessary for carrying a pregnancy and the amount of oxygen supplied to the fetus. It results in such problems as fetal growth retardation, preterm labor, and congenital abnormalities. The purpose of the study is to evaluate the impact of SHS on the fetoplacental unit and neonatal outcome via a comparative cross-sectional survey of 50 pregnant women with and without smoke exposure. The clinical, laboratory, and statistical analysis assesses maternal and fetal health consequences, with focused attention being directed at placental performance and the initial phase of newborn growth. The results for patients will help in developing preventive strategies, and broaden the current knowledge about sociogenesis and clinical relevance of passive smoking during pregnancy and subsequent directions in the intervention.

KEYWORDS: Passive smoking, pregnancy outcomes, neonatal health, placental function, maternal health.

Introduction

Research has established that the emission of tobacco smoke has many chemicals that act as a reproductive hazard. Environmental tobacco smoke or passive smoking whereby an individual is exposed to smoke they did not generate has been found to cause severe gestational and neonatal effects. Carbon monoxide hinders fetal oxygen transport because it bonds with hemoglobin; nicotine complicates pregnancy by inhibiting the secretion of pregnancy hormones, for example, progesterone and prolactin. This hormonal change is not favorable for the development of the placenta and raises the chances of impossibility of implantation, premature labor, or placental problems. Epidemiological research confirmed the dose-response relationship between TSB and pregnancy outcomes such as low birth weight, congenital abnormalities, and delayed

developmental development problems in newborns. The offspring born from mothers who are passively exposed to tobacco smoke are also more prone to contract respiratory and cardiovascular illnesses. The purpose of this research shall be to examine the impact of passive smoking on the progress of pregnancy and the newborn. Questionnaire-integrated clinical, psychological, and laboratory interventions to determine the sociological factors of passive smoking and its effects on maternal and neonatal health, will be employed. The findings should have the potential to refine the preventive approaches and optimize the perinatal preferences of women.

A large number of studies describe how maternal and fetal health can be harmed by tobacco smoke. The components of tobacco smoke include carbon monoxide, nicotine, heavy metals, and carcinogenic compounds that limit physiological activities critical to a healthy pregnancy. Literature suggests that passive smoking raises pregnancy risks including low birth weight, fetal hypoxia, and congenital anomalies. Martin, the leading toxicant in tobacco smoke, has a higher affinity to fetal hemoglobin than oxygen and thus impairs oxygen delivery causing a chronic hypoxic state. Semenova et al. (2016) concluded that this causes poor fetal growth and development. Just like estrogen and progesterone, nicotine inhibits the secretion of essential pregnancy hormones affecting placental development and contributing to increased rates of miscarriages, placental abruption, and preterm labor. The author found in epidemiological studies that there is a direct relationship between passive smoking of women and adverse outcomes including placental previa and perinatal mortality. However, cross-sectional health effects associated with children born to mothers who had been exposed to tobacco smoke include respiratory and cardiovascular diseases in the long run. There have also been the sociogenesis of passive smoking studies done in the recent past where most researchers have evaluated factors that have psychological and or socio-economic impacts on exposure levels. Smoking cessation and second-hand smoking exposure prevention are contingent on the medical and necessity of maternal and neonatal health.

In this present research, a systematic method of working is used to analyze the impacts of passive smoking on pregnancy, labor, and neonatal factors. A comparative observational design is used, involving 50 pregnant women categorized into two groups: one which comprises healthy pregnant women who have not been exposed to tobacco smoke during pregnancy, and the other which consists of women who became passive smokers during pregnancy. Methods of data collection encompass clinical assessment, laboratory tests, and psychological tests as well. Clinical assessments examine maternity concerns and pregnancy development, and the children's birth, preterm birth factors, and neonatal status by analyzing placental features, birth weights, and other newborn findings. Laboratory analyses consist in evaluating maternal blood and umbilical cord blood for indexes that are carbon monoxide and cotinine, a nicotine derivative. It is desired that these analyses provide the comprehension of the biochemical impacts of passive smoking on the fetoplacental system and neonatal health. Sample surveys are employed by questionnaires to determine sociological and psychological determinants of passive smoking exposure. They include such factors as lifestyle practices, socio-economic status, and relative awareness of the dangers of passive smoking. The data collected is analyzed by using statistical techniques. Measures of descriptive statistics bring an overview of the findings and inferential statistics like regression analysis determine the relationship of the extent of passive smoking exposure with adverse maternal and neonatal outcomes.

However, practices the highest ethical consideration in the study, is respect for participant's privacy and their consent to participate in the study. The present cross-sectional methodological design will give a vast picture of the effects of passive smoking on maternal and neonatal health. It also aims to reveal sociological trends and clinical applications and make positive contributions to the formulation of efficient prevention and accurate intervention programs. The study evaluated 50 pregnant women, divided into two groups: one comprised of women who were exposed to passive smoking during pregnancy and a comparison group with no exposure. The findings revealed significant differences in maternal, placental, and neonatal health outcomes between the groups:

1. Maternal Health:

- Women exposed to passive smoking exhibited higher rates of pregnancy complications, including placental abnormalities (e.g., previa and abruption) and preterm labor.
- Biochemical analysis indicated elevated levels of carbon monoxide and cotinine in the blood, correlating with reduced oxygen transport and hormonal imbalances.

2. Placental Function:

- Placental samples from the exposed group showed signs of reduced thickness and functionality, with impaired blood flow and chronic hypoxia.
- Histological analysis indicated increased levels of oxidative stress markers, contributing to placental insufficiency.

3. Neonatal Outcomes:

- Neonates in the exposed group had lower birth weights (average 2.8 kg vs. 3.2 kg in the control group) and higher incidences of congenital anomalies and early neonatal complications.
- Apgar scores in the first and fifth minutes were significantly lower among neonates from the exposed group.

In light of these, the findings affirm the ill-health consequences of passive smoking on pregnancy and neonate wellbeing. Carbon monoxide has a high affinity for hemoglobin, thus organizing a low level of oxygen in the fetus leading to a permanent hypoxic state and developmental retardation. Nicotine also interferes with hormonal influence preventing placenta growth and normal fetal organ development. Reduced placental size and abnormal placental location in the exposed group are consistent with previous findings and the reasons behind the elevated risk of previa and abruption because of poor perfusion and placental villous trophoblast modulation. Such conditions predispose the women to premature labor and neonatal poor prognosis characterized by low birth weight and poor Apgar scores. Furthermore, the observed association between passive smoking exposure and raised oxidative stress indicates dire early intercessions. Thus, sociological material indicates that female patients in the exposed group were less informed as to the hazards of passive smoking, which underlines the need for specialized educational campaigns.

Conclusion

This paper establishes the threat of passive smoking on maternal, placental, and neonatal outcomes. The evidence presented herein clearly shows that maternal tobacco smoke exposure during pregnancy raises the risks of such pregnancy complications as placental abnormalities, preterm labor, and neonatal adverse outcomes including LBW and congenital anomalies. The former harms hormonal regulation for pregnancy impairs the function of the placenta, and reduces the oxygen supply to the fetus resulting in chronic hypoxia and developmental delays. This study realizes the importance of education and preventive measures in controlling the effects of passive smoking. The results of questionnaires used in the sociological study show that many pregnant women lack adequate knowledge of the adverse effects of second-hand smoke; hence the need to employ intervention strategies. With prevention and education on changes that would surround passive smoking during pregnancy it will be possible to decrease the cases and at the same time have healthy birthing mothers and babies. Such outcomes are useful for extending knowledge on the components of passive smoking and its clinical and sociological penalties. They also offer the basis for soon formulating practical evidence-based policies and preventive measures for reducing exposure to tobacco smoke during pregnancy. More attention should be paid to the further study of what passive smoking does to the development of children and how particular strategies can be improved to produce healthier mothers and their offspring.

REFERENCES

1. Семенова, Т. В., Аржанова, О. Н., Беспалова, О. Н., & Прокопенко, В. М. (2016). Особенности течения беременности и исходов родов при табакокурении. *Акушерство и гинекология*. Москва: ГЭОТАР-Медиа.
2. Табакокурение как фактор риска болезней человека; Рекомендации для беременных женщин и молодых матерей. Консультационный этап - Программы выхода беременности и общественное здравоохранение; Rivers & Associates. (2005). Виктория, ВС, с. 24.
3. Сахарова, Г. М., Антонов, Н. С., & Андреева, С. А. (2007). Табакокурение как фактор риска болезней человека. *Респираторная медицина*. Москва: Издательство ГЭОТАР–Медиа.
4. Умаров, Б. Х., Хусонов, К. С., & Назаров, Ф. Т. (2018). Тамаки тутуни таъсири ва репродуктив саломатлик. *Ўзбекистон тиббиёт журнали*, 2(3), 45-52.
5. Мендола, П., Ха, С., & Поллак, А. З. (2015). Загрязнение воздуха и преждевременные роды: последние находки и направления для будущих исследований. *Current Epidemiology Reports*, 2(3), 156-162. <https://doi.org/10.1007/s40471-015-0041-4>.
6. Султонов, А. Х. (2020). Тамаки тутунининг оналар ва болалар соғлиғига таъсири: ижтимоий ва клиник тадқиқотлар. *Тиббиётда инновациялар журнали*, 1(4), 67-74.
7. Семенова, Т. В., Аржанова, О. Н., Беспалова, О. Н., & Прокопенко, В. М. (2016). Особенности течения беременности и исходов родов при табакокурении. *Акушерство и гинекология*. Москва: ГЭОТАР-Медиа.
8. Johansson, A. K., Pliadou, A., Bergvall, N., Tuvemo, T., Norman, M., & Cnattingius, S. (2008). Risk of reduced intellectual performance in male adolescents exposed to maternal smoking during pregnancy. *Pediatric Research*, 64(1), 90-94. <https://doi.org/10.1203/PDR.0b013e318176bc48>.