



# Innovative Approaches to Ensuring the Quality and Safety of Functional Confectionery Products Based on Flour

**M. Z. Ashurova**

Bukhara state technical university, docent of the Department of "Food Technologies and Service"

**A. Xayrullayev**

Bukhara state technical university, Master's degree in "Food Safety"

## Annotation

The growing consumer demand for healthier confectionery alternatives has accelerated the development of functional flour-based products enriched with bioactive compounds. This review examines innovative approaches to ensuring quality and safety in functional confectionery production, focusing on flour modification techniques, biofortification strategies, and quality control systems. Current methodologies including high-performance liquid chromatography (HPLC), near-infrared spectroscopy (NIRS), and advanced microbiological testing are analyzed. The integration of HACCP principles with novel preservation technologies such as microencapsulation and natural antimicrobials demonstrates significant potential for extending shelf life while maintaining nutritional value. Findings indicate that combined approaches involving modified atmosphere packaging, optimal water activity control (0.3-0.6 aw), and strategic incorporation of functional ingredients can enhance both safety parameters and consumer acceptance. Future developments should prioritize sustainable production methods and personalized nutrition approaches to meet evolving market demands.

**Keywords:** functional confectionery, flour-based products, quality assurance, food safety, biofortification, HACCP.



This is an open-access article under the [CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/) license

## INTRODUCTION

The global confectionery market has witnessed a paradigm shift toward functional food products that provide health benefits beyond basic nutrition. Flour-based confectionery products, including cookies, wafers, cakes, and pastries, represent a significant segment of the food industry, with annual production exceeding 50 million tons worldwide. However, traditional confectionery products are often associated with high caloric content, refined sugars, and limited nutritional value, contributing to increasing rates of obesity, diabetes, and cardiovascular diseases.

Functional confectionery products emerge as a promising solution, incorporating bioactive compounds such as dietary fibers, antioxidants, vitamins, minerals, and probiotics into conventional formulations. The functional food market has experienced exponential growth, with projections indicating a compound annual growth rate (CAGR) of 8.5% through 2028. This trend reflects consumer awareness of diet-related health issues and the desire for convenient, health-promoting food options.


### **QUALITY AND SAFETY CHALLENGES**

Despite the promising potential, functional confectionery products face unique quality and safety challenges. The incorporation of functional ingredients often affects sensory attributes, texture, shelf stability, and microbiological safety. Flour-based matrices are susceptible to lipid oxidation, moisture migration, microbial contamination, and mycotoxin formation during storage. Furthermore, the interaction between functional ingredients and flour components can lead to unpredictable changes in product characteristics.

Ensuring consistent quality and safety requires innovative approaches that address these multifaceted challenges while maintaining consumer acceptability and nutritional efficacy. Traditional quality control methods may prove insufficient for complex functional formulations, necessitating the integration of advanced analytical techniques and predictive modeling.

### **RESEARCH OBJECTIVES**

This review aims to synthesize current knowledge on innovative approaches to ensuring quality and safety in functional flour-based confectionery products.



**Picture 1. Specific objectives include**

### **METHODS**

*Literature Search Strategy.* A comprehensive literature review was conducted using multiple scientific databases including PubMed, Scopus, Web of Science, and Google Scholar. The search strategy employed Boolean operators combining keywords: "functional confectionery," "flour-based products," "quality control," "food safety," "biofortification," "HACCP," "preservation technologies," and "analytical methods." The search covered publications from 2015 to 2025, prioritizing peer-reviewed research articles, systematic reviews, and authoritative industry reports.

*Inclusion and Exclusion Criteria.* Studies were included if they focused on:


- ✓ Quality assessment methodologies for flour-based confectionery products

- ✓ Safety assurance systems and hazard analysis
- ✓ Functional ingredient incorporation and stability
- ✓ Novel preservation and packaging technologies
- ✓ Analytical techniques for quality parameter determination

Exclusion criteria eliminated studies focusing solely on non-flour confectionery, purely theoretical modeling without experimental validation, and publications without quality peer review.

*Data Extraction and Synthesis.* Relevant information was systematically extracted including study design, methodological approaches, key findings, and practical applications. Data synthesis employed a narrative review approach, categorizing information into thematic areas: ingredient innovation, quality control technologies, safety management systems, and preservation strategies. Critical analysis focused on identifying knowledge gaps, methodological limitations, and emerging trends.

*Analytical Framework.* The review employed a multi-dimensional analytical framework encompassing:



## RESULTS

Heat-moisture treatment (HMT) and annealing have emerged as effective physical modification techniques for improving flour functionality in confectionery applications. Research demonstrates that HMT conducted at 100-120°C with 20-30% moisture content for 1-16 hours significantly alters starch gelatinization properties, enhancing thermal stability and reducing digestibility. Studies report that HMT-modified wheat flour exhibits 15-25% reduction in rapidly digestible starch, making it suitable for low-glycemic functional products.

Ultrasonic treatment (20-40 kHz, 400-600 W) represents another promising approach, inducing controlled structural modifications without chemical additives. Ultrasound-treated flour demonstrates improved water absorption capacity (8-12% increase), reduced dough development time (20-30% decrease), and enhanced gluten network formation, resulting in superior textural properties in baked confectionery products.

Enzymatic modification using transglutaminase, glucose oxidase, and xylanase enables targeted functional improvements. Transglutaminase application (0.1-1.0% flour weight) creates cross-linked protein networks, improving dough strength and product structure. Studies indicate that

enzymatically modified flour produces confectionery products with 20-35% improved volume retention and extended shelf life (15-20 days increase at ambient storage).

**BIOFORTIFICATION STRATEGIES.** Biofortification through ingredient substitution has gained substantial research attention. Partial wheat flour replacement with nutrient-dense alternatives including:

- Whole grain flours: Oat, barley, rye (10-30% substitution) increasing dietary fiber by 40-60%
- Pseudocereal flours: Quinoa, amaranth, buckwheat (15-25% substitution) enhancing protein quality and mineral content
- Legume flours: Chickpea, lentil, soybean (10-20% substitution) improving protein content by 25-45%
- Functional flours: Sprouted, germinated, fermented variants providing enhanced bioavailability

Research demonstrates that 20% oat flour incorporation increases  $\beta$ -glucan content to 2.5-3.5 g per 100g serving, supporting cardiovascular health claims. Similarly, 15% chickpea flour addition elevates protein content by 30-40% while providing resistant starch and prebiotic effects.

Near-infrared spectroscopy (NIRS) has revolutionized rapid, non-destructive quality assessment in flour-based confectionery production. NIRS calibration models achieve prediction accuracy ( $R^2 > 0.95$ ) for critical parameters including moisture content, protein levels, fat composition, and ash content. Portable NIRS devices enable real-time in-line monitoring, reducing analysis time from hours to seconds while eliminating sample preparation requirements.

Fourier-transform infrared spectroscopy (FTIR) provides detailed molecular fingerprinting for authentication and quality verification. FTIR successfully detects adulterations, monitors lipid oxidation through carbonyl group formation, and tracks protein structural changes during processing. Studies report FTIR classification accuracy exceeding 95% for flour variety identification and contamination detection.

High-performance liquid chromatography (HPLC) remains the gold standard for quantifying bioactive compounds, vitamins, antioxidants, and potential contaminants. Advanced HPLC methodologies coupled with mass spectrometry (HPLC-MS/MS) enable simultaneous multi-residue analysis with detection limits in the parts-per-billion range.

**Recent applications include:**


- ✓ Polyphenol profiling in enriched formulations (LOD: 0.05-0.5  $\mu\text{g/g}$ )
- ✓ Mycotoxin screening (aflatoxins, ochratoxin A, deoxynivalenol)
- ✓ Acrylamide monitoring in baked products
- ✓ Vitamin stability assessment during storage

Gas chromatography-mass spectrometry (GC-MS) facilitates volatile compound analysis, identifying off-flavors, oxidation products, and aroma profiles. Electronic nose technology, utilizing sensor arrays mimicking human olfaction, provides rapid quality classification with 85-95% accuracy compared to trained sensory panels.

Instrumental texture analysis using texture analyzers provides objective quantification of mechanical properties critical to consumer acceptance. Key parameters including hardness, fracturability, chewiness, and cohesiveness correlate strongly with sensory perception ( $r = 0.75-0.92$ ). Dynamic mechanical analysis (DMA) and oscillatory rheology characterize viscoelastic behavior during processing, enabling formulation optimization.

Advanced imaging technologies including X-ray microtomography and confocal laser scanning microscopy reveal microstructural characteristics influencing texture and stability. Three-dimensional porosity analysis predicts moisture migration patterns and oxidative stability with improved accuracy.

HACCP Implementation. Hazard Analysis and Critical Control Points (HACCP) implementation has become mandatory in functional confectionery production.



**Picture 2. Critical control points**

Studies demonstrate that comprehensive HACCP implementation reduces food safety incidents by 60-75% and improves regulatory compliance. Integration with automated monitoring systems enables real-time hazard detection and corrective action implementation.

Advanced microbiological testing methods accelerate pathogen detection while improving sensitivity. Polymerase chain reaction (PCR) and loop-mediated isothermal amplification (LAMP) reduce detection time from 3-5 days to 2-6 hours for pathogens including Salmonella, E. coli, and Listeria. Flow cytometry enables rapid total viable count determination within 30 minutes, facilitating immediate production decisions.

Predictive microbiology models estimate microbial growth kinetics under various storage conditions. Validated mathematical models (Gompertz, Baranyi) predict shelf life with 90-95% accuracy, considering temperature fluctuations, water activity, and pH interactions.

Allergen management represents a critical safety concern in functional confectionery. Implementation of dedicated production lines, thorough cleaning validation (ATP bioluminescence testing), and sensitive detection methods (ELISA, lateral flow devices) minimize

cross-contact risks. Quantitative allergen testing achieves detection limits of 0.5-5 ppm, ensuring compliance with labeling regulations.

Natural antimicrobial incorporation provides clean-label preservation alternatives. Effective compounds include:

- Essential oils: Oregano, thyme, cinnamon (0.1-0.5%) exhibiting broad-spectrum antimicrobial activity
- Plant extracts: Green tea, rosemary, grape seed polyphenols (0.02-0.1%) providing antioxidant and antimicrobial effects
- Organic acids: Citric, ascorbic, sorbic acids optimizing pH and inhibiting microbial growth
- Bacteriocins: Nisin, natamycin targeting specific spoilage organisms

Research indicates that synergistic combinations (hurdle technology) achieve microbial inhibition equivalent to synthetic preservatives while maintaining sensory acceptability. Microencapsulation of volatile antimicrobials in cyclodextrin, liposomes, or polysaccharide matrices controls release kinetics and minimizes organoleptic impact.

Modified atmosphere packaging (MAP) extends shelf life by modifying gas composition surrounding the product. Optimal gas mixtures for flour-based confectionery typically include:

- Low oxygen (0-5%) inhibiting lipid oxidation and aerobic spoilage
- Elevated carbon dioxide (20-60%) providing fungistatic effects
- Nitrogen (balance) preventing package collapse

Studies demonstrate that MAP combined with oxygen scavengers extends shelf life by 50-150% compared to conventional packaging. Active packaging systems incorporating ethanol emitters, moisture regulators, and antimicrobial films provide additional protection.

Water activity (aw) control represents the most critical factor influencing microbial stability and chemical reactions. Optimal aw ranges for flour-based confectionery (0.3-0.6) prevent microbial growth while minimizing lipid oxidation. Precise aw control achieved through:

- Humectant selection (glycerol, sorbitol, propylene glycol)
- Formulation optimization balancing ingredients
- Processing conditions (baking time/temperature)
- Packaging barrier properties

Predictive models correlate aw with glass transition temperature, enabling shelf-life optimization across storage conditions.

The findings demonstrate that effective quality and safety assurance in functional confectionery requires integrated, multi-factorial approaches rather than isolated interventions. Synergistic combinations of flour modification, biofortification, advanced analytics, and preservation technologies yield optimal results. For instance, heat-moisture treated flour enriched with antioxidant compounds, packaged under modified atmosphere, and monitored using NIRS-based quality control achieves superior nutritional value, extended shelf life, and consistent quality compared to single-approach strategies.

The implementation of predictive modeling throughout the production chain enables proactive quality management. Mathematical models integrating ingredient properties, processing parameters, and environmental conditions accurately predict product characteristics, facilitating

formulation optimization before physical production. This approach reduces development time by 40-60% and minimizes material waste.

Despite significant advances, several challenges persist. Functional ingredient incorporation often compromises sensory attributes, particularly texture and flavor. Consumer acceptance studies indicate that sensory quality remains the primary purchase driver, superseding health benefits. Balancing nutritional enhancement with palatability requires sophisticated formulation strategies and potentially consumer education initiatives.

Cost implications represent another significant barrier. Advanced analytical equipment (HPLC-MS, NIRS), modified packaging systems, and premium functional ingredients increase production costs by 25-50%. Market viability depends on consumer willingness to pay premium prices, which varies substantially across demographics and regions.

Regulatory complexity poses additional challenges, particularly for novel functional ingredients and health claims. Substantiation requirements for functional claims necessitate extensive clinical evidence, representing substantial time and financial investments. Harmonization of international regulations would facilitate market access and encourage innovation.

Sustainable production practices are increasingly prioritized by consumers, retailers, and regulators. Functional confectionery development should incorporate sustainability principles including:

- ✓ Utilization of underutilized crops and by-products (bran, pulses, ancient grains)
- ✓ Energy-efficient processing technologies
- ✓ Biodegradable or recyclable packaging materials
- ✓ Local ingredient sourcing reducing transportation impact
- ✓ Waste minimization through ingredient optimization

Research demonstrates that sustainable practices can enhance brand perception and consumer loyalty while potentially reducing operational costs through improved efficiency.

**Personalized Nutrition:** Advances in nutrigenomics and consumer health monitoring enable personalized product formulations addressing individual nutritional needs, genetic predispositions, and health goals. Digital platforms facilitating customized confectionery production represent a promising market opportunity.

**Artificial Intelligence Integration:** Machine learning algorithms analyzing vast datasets from production, quality control, and consumer feedback enable continuous optimization and predictive maintenance. AI-driven formulation tools accelerate product development and improve success rates.

**Novel Functional Ingredients:** Ongoing research into bioactive compounds from marine sources, microalgae, insects, and cellular agriculture expands functional ingredient options. These emerging ingredients offer superior nutritional profiles and sustainability advantages.

**Clean Label Movement:** Consumer demand for transparent, minimally processed products with recognizable ingredients intensifies. Future innovations must balance technological sophistication with clean label positioning through natural alternatives and processing optimization.

**Blockchain Technology:** Implementation of blockchain-based traceability systems enhances transparency, facilitates rapid recall execution, and builds consumer trust. Complete supply chain visibility from farm to consumer becomes increasingly feasible.

## CONCLUSION

Ensuring quality and safety in functional flour-based confectionery products requires comprehensive, integrated approaches combining ingredient innovation, advanced analytical methods, robust safety systems, and effective preservation technologies. Current research demonstrates substantial progress in flour modification techniques, biofortification strategies, and quality control methodologies. The successful implementation of HACCP principles integrated with predictive modeling and advanced analytics significantly enhances product safety and consistency.

Future developments should prioritize sustainable production methods, personalized nutrition approaches, and clean label formulations while maintaining sensory acceptability and economic viability. Continued interdisciplinary research incorporating food science, nutrition, engineering, and consumer behavior will drive innovation in this dynamic field. The functional confectionery sector represents a promising opportunity to deliver health benefits through convenient, enjoyable food products, contributing to public health improvement while meeting evolving consumer demands.

## REFERENCES

1. Regulation, E. C. (2006). Regulation (EC) No 1924/2006 of the European Parliament and of the Council on nutrition and health claims made on foods. *\*Official Journal of the European Union\**, 404, 9-25.
2. Bigliardi, B., & Galati, F. (2013). Innovation trends in the food industry: the case of functional foods. *\*Trends in Food Science & Technology\**, 31(2), 118-129.
3. Gul, K., Yousuf, B., Singh, A. K., Singh, P., & Wani, A. A. (2015). Rice bran: Nutritional values and its emerging potential for development of functional food—A review. *\*Bioactive Carbohydrates and Dietary Fibre\**, 6(1), 24-30.
4. Ktenioudaki, A., & Gallagher, E. (2012). Recent advances in the development of high-fibre baked products. *\*Trends in Food Science & Technology\**, 28(1), 4-14.
5. Pasqualone, A., Delvecchio, L. N., Gambacorta, G., Laddomada, B., Urso, E., Mazzaglia, A., ... & Summo, C. (2016). Effect of supplementation with wheat bran aqueous extracts obtained by ultrasound-assisted technologies on the sensory properties and the antioxidant activity of dry pasta. *\*Natural Product Research\**, 30(20), 2303-2309.
6. Rosell, C. M., Rojas, J. A., & Benedito de Barber, C. (2001). Influence of hydrocolloids on dough rheology and bread quality. *\*Food Hydrocolloids\**, 15(1), 75-81.
7. Struck, S., Plaza, M., Turner, C., & Rohm, H. (2016). Berry pomace—a review of processing and chemical analysis of its polyphenols. *\*International Journal of Food Science & Technology\**, 51(6), 1305-1318.
8. Gómez, M., Doyagüe, M. J., & De la Hera, E. (2012). Addition of pin-milled pea flour and air-classified fractions in layer and sponge cakes. *\*LWT-Food Science and Technology\**, 46(1), 142-147.
9. Poutanen, K., Sozer, N., & Della Valle, G. (2014). How can technology help to deliver more of grain in cereal foods for a healthy diet? *\*Journal of Cereal Science\**, 59(3), 327-336.
10. Mishra, S., Monro, J., & Hedderley, D. (2008). Effect of processing on slowly digestible starch and resistant starch in potato. *\*Starch-Stärke\**, 60(9), 500-507.